arXiv:1202.2278v1 [physics.gen-ph] 9 Feb 2012

QCD ghost $f(T)$-gravity model

K. Karami[1,2*], A. Abdolmaleki[1†]

[1]Department of Physics, University of Kurdistan, Pasdaran St., Sanandaj, Iran
[2]Research Institute for Astronomy and Astrophysics of Maragha (RIAAM), Maragha, Iran

February 13, 2012

Abstract

We reconstruct a $f(T)$ modified teleparallel gravity model corresponding to the ghost dark energy scenario. We calculate the effective torsion equation of state and deceleration parameters of the ghost $f(T)$-gravity model. We conclude that the equation of state parameter behaves like a freezing quintessence model. Furthermore, the deceleration parameter can justify the transition from the cosmic deceleration to acceleration phase of the universe. Also the present value of the deceleration parameter is in good agreement with the recent observations.

PACS numbers: 95.36.+x, 04.50.Kd
Keywords: Dark energy, Modified theories of gravity

1 Introduction

Here, our aim is to reconstruct a $f(T)$ modified teleparallel gravity model according to the ghost dark energy (GDE) scenario.

The action of $f(T)$-gravity is given by [1]-[10]

$$I = \frac{1}{2k^2}\int \mathrm{d}^4x\ e\ \Big[f(T) + L_m\Big], \tag{1}$$

where $k^2 = 8\pi G$, $e = \det(e^i_\mu)$ and e^i_μ is the vierbein field which is used as a dynamical object in teleparallel gravity. Also T and L_m are the torsion scalar and the Lagrangian density of the matter inside the universe, respectively.

Taking the variation of the action (1) with respect to the vierbein e^i_μ, the modified Friedmann equations in the spatially flat Friedmann-Robertson-Walker (FRW) universe can be obtained in the standard forms

$$\frac{3}{k^2}H^2 = \rho_m + \rho_T, \tag{2}$$

$$\frac{1}{k^2}(2\dot{H} + 3H^2) = -(p_m + p_T), \tag{3}$$

where

$$\rho_T = \frac{1}{2k^2}(2Tf_T - f - T), \tag{4}$$

---

[*] KKarami@uok.ac.ir
[†] AAbdolmaleki@uok.ac.ir

$$p_T = -\frac{1}{2k^2}[-8\dot{H}Tf_{TT} + (2T - 4\dot{H})f_T - f + 4\dot{H} - T], \tag{5}$$

and

$$T = -6H^2. \tag{6}$$

Here, $H = \dot{a}/a$ is the Hubble parameter and the subscript T in f_T denotes a derivative with respect to the torsion scalar T. Also ρ_m and p_m are the energy density and pressure of the matter inside the universe, respectively. Furthermore, ρ_T and p_T are the torsion contributions to the energy density and pressure, respectively.

The energy conservation equations are still given by

$$\dot{\rho}_m + 3H(\rho_m + p_m) = 0, \tag{7}$$

$$\dot{\rho}_T + 3H(\rho_T + p_T) = 0. \tag{8}$$

The effective torsion equation of state (EoS) parameter is defined as [5, 7]

$$\omega_T = \frac{p_T}{\rho_T} = -1 - \frac{\dot{T}}{3H}\left(\frac{2Tf_{TT} + f_T - 1}{2Tf_T - f - T}\right). \tag{9}$$

With the help of Eqs. (2), (4) and (6) one can get

$$\rho_m = \frac{1}{16\pi G}(f - 2Tf_T). \tag{10}$$

For the pressureless matter, i.e. $p_m = 0$, from Eqs. (2) to (5) one can obtain

$$\dot{H} = -\frac{4\pi G\rho_m}{f_T + 2Tf_{TT}}. \tag{11}$$

Inserting Eq. (10) into (11) and using $\dot{T} = -12H\dot{H}$ gives

$$\dot{T} = 3H\left(\frac{f - 2Tf_T}{f_T + 2Tf_{TT}}\right). \tag{12}$$

Using the above relation, the effective EoS parameter (9) yields

$$\omega_T = -\frac{f/T - f_T + 2Tf_{TT}}{(f_T + 2Tf_{TT})(f/T - 2f_T + 1)}. \tag{13}$$

Here, we also calculate the deceleration parameter

$$q = -1 - \frac{\dot{H}}{H^2}, \tag{14}$$

which can be compared with the observations. Using Eqs. (6) and (12) the deceleration parameter (14) leads to

$$q = 2\left(\frac{f_T - Tf_{TT} - \frac{3f}{4T}}{f_T + 2Tf_{TT}}\right). \tag{15}$$

2 Ghost $f(T)$-gravity model

The dark torsion contribution in $f(T)$-gravity can justify the observed acceleration of the universe without resorting to the DE. This motivates us to reconstruct a $f(T)$-gravity model according to the GDE model which has been recently motivated from the Veneziano ghost of quantum choromodynamics (QCD). The GDE density is given by [11]

$$\rho_D = \alpha H, \tag{16}$$

where α is a constant with dimension [energy]^3. Several aspects of this new paradigm have been investigated in the literature [12].

With the help of Eq. (6) one can rewrite (16) as

$$\rho_D = \alpha \left(-\frac{T}{6}\right)^{1/2}. \tag{17}$$

Equating (4) and (17), i.e. $\rho_T = \rho_D$, we obtain the following differential equation

$$2Tf_T - f - T - \beta\sqrt{-T} = 0, \tag{18}$$

where

$$\beta = \frac{2k^2\alpha}{\sqrt{6}}. \tag{19}$$

Solving Eq. (18) yields the $f(T)$-gravity corresponding to the GDE model as

$$f(T) = T + \sqrt{-T}\left(\epsilon + \frac{\beta}{2}\ln(-T)\right), \tag{20}$$

where ϵ is an integration constant that can be determined from a boundary condition. Following [13] to recover the present day value of Newtonian gravitational constant we need to have

$$f_T(T_0) = 1, \tag{21}$$

where $T_0 = -6H_0^2$ is the torsion scalar at the present time. Applying the above boundary condition to the solution (20) one can obtain

$$\epsilon = \beta\left(1 - \frac{1}{2}\ln(-T_0)\right). \tag{22}$$

Substituting this into Eq. (20) gives

$$f(T) = T + \beta\sqrt{-T}\left[\frac{1}{2}\ln\left(\frac{T}{T_0}\right) - 1\right]. \tag{23}$$

The parameter β can be obtained by inserting Eq. (23) into the modified Friedmann equation (2). Solving the resulting equation for the present time gives

$$\beta = \sqrt{6}H_0(1 - \Omega_{m_0}), \tag{24}$$

where $\Omega_{m_0} = \frac{k^2\rho_{m_0}}{3H_0^2}$ is the dimensionless matter energy density and the index 0 denotes the value of a quantity at present. The evolution of the ghost $f(T)$-gravity model, Eq. (23), versus T/T_0 is plotted in Fig. 1. It shows that the magnitude of ghost $f(T)$ model increases when T/T_0 increases. At present time $T/T_0 = 1$ we have $f(T_0) = -7974$.

Note that the ghost $f(T)$-gravity (23) satisfies the condition

$$\lim_{T\to\infty} f/T \to 1, \tag{25}$$

at high redshift which is compatible with the primordial nucleosynthesis and CMB constraints [6, 8].

Inserting Eq. (23) into (12) and using $H = (-T/6)^{1/2}$ yields

$$t - t_i = \frac{1}{\sqrt{6}} \int_{T_{\rm i}}^{T} \left(\frac{2\sqrt{-T} - \beta}{\beta\sqrt{-T} + T} \right) \frac{{\rm d}T}{T}. \tag{26}$$

Integrating the above relation analytically gives

$$t = \sqrt{\frac{2}{3}} \left[\frac{1}{\sqrt{-T}} + \frac{1}{\beta} \ln \left(\frac{\sqrt{-T}}{\sqrt{-T} - \beta} \right) \right], \quad \sqrt{-T} \geq \beta, \tag{27}$$

where we take $T_i = -6H_i^2 = -\infty$ at $t_i = 0$. Note that the condition $\sqrt{-T} \geq \beta$ is necessary due to having a real time. Using Eq. (24) the condition $\sqrt{-T} \geq \beta$ can be rewritten as $T/T_0 \geq (1 - \Omega_{m_0})^2$. Figure 2 shows the time evolution of the fractional torsion scalar T/T_0. It clears that T/T_0 decreases with increasing the time. Figure 2 illustrates that at early $(t/t_0 \to 0)$ and late $(t/t_0 \to \infty)$ times we have $T/T_0 \to \infty$ and $T/T_0 \to (1 - \Omega_{m_0})^2 = 0.55$, respectively, where we take $\Omega_{m_0} = 0.26$ [13].

Inserting Eq. (23) into (13) gives the effective torsion EoS parameter of the ghost $f(T)$-gravity model as

$$\omega_T = -\frac{T}{2T + \beta\sqrt{-T}}. \tag{28}$$

The time evolution of the EoS parameter (28) is plotted in Fig. 3. It shows that at early times $(t/t_0 \to 0)$ we have $\omega_T \to -0.5$ and at late times $(t/t_0 \to \infty)$ we get $\omega_T \to -1$ which acts like ΛCDM. Also at present time we have $\omega_{T_0} = -0.79$. Figure 3 clears that the torsion EoS parameter of the ghost $f(T)$-gravity model behaves like freezing quintessence DE [14] and cannot cross the phantom divide line. This result is in complete agreement with that obtained for the GDE model in the Einstein gravity [11, 12].

Inserting Eq. (23) into (15) one can obtain the deceleration parameter

$$q = \frac{T + 2\beta\sqrt{-T}}{2T + \beta\sqrt{-T}}. \tag{29}$$

Figure 4 shows the time evolution of the deceleration parameter (29). It clears that at early times $(t/t_0 \to 0)$ we have $q \to 0.5$ which corresponds to the matter dominated universe. Also at late times $(t/t_0 \to \infty)$ we get $q \to -1$ which behaves like the de Sitter universe. Figure 4 illustrates that at $t/t_0 = 0.57$ we have a cosmic deceleration $q > 0$ to acceleration $q < 0$ transition which is compatible with the observations [15]. Also at present time we get $q_0 \simeq -0.4$ which is in good agreement with recent observational results $-1.4 \leq q_0 \leq -0.3$ [15].

3 Conclusions

Here, we established a correspondence between the GDE model and $f(T)$ modified teleparallel gravity. In the framework of $f(T)$ theory, we considered a spatially flat FRW universe filled only with ordinary matter. Then, we reconstructed a $f(T)$ model according to the GDE scenario. We concluded that the condition $f/T \to 1$ is satisfied at high redshift ($T \to \infty$) which is compatible with the primordial nucleosynthesis and CMB constraints. We also obtained the time evolution of the dark torsion scalar T. Furthermore, we calculated the effective EoS and deceleration parameters of the ghost $f(T)$-gravity model. Our results show the following.

(i) The EoS parameter ω_T varies from -0.5 at early times to -1 at late times, which is similar to freezing quintessence type of DE. For the present time we obtain $\omega_{T_0} = -0.79$.

(ii) The deceleration parameter q at early and late times behaves like the matter dominated and de Sitter universe, respectively. Also at near past $t/t_0 = 0.57$ we have a cosmic deceleration ($q > 0$) to acceleration ($q < 0$) transition. The deceleration parameter $q_0 \simeq -0.4$ at the present time is compatible with the recent observations.

All in all, we pointed out that the dynamics of the universe described by the GDE in the Einstein gravity can be explained by $f(T)$ theory without resorting to the DE paradigm.

Acknowledgements

The work of K. Karami has been supported financially by Research Institute for Astronomy & Astrophysics of Maragha (RIAAM), Maragha, Iran.

References

[1] G.R. Bengochea, R. Ferraro, Phys. Rev. D **79**, 124019 (2009);
G.R. Bengochea, Phys. Lett. B **695**, 405 (2011).

[2] R.J. Yang, Eur. Phys. J. C **71**, 1797 (2011);
S.C. Ulhoa, J.F. da Rocha Neto, J.W. Maluf, Int. J. Mod. Phys. D **19**, 1925 (2010);
K. Bamba, C.Q. Geng, C.C. Lee, arXiv:1008.4036;
K. Bamba, et al., JCAP **01**, 021 (2011);
R. Zheng, Q.G. Huang, JCAP **03**, 002 (2011).

[3] K.K. Yerzhanov, et al., arXiv:1006.3879;
R. Myrzakulov, arXiv:1008.4486;
R. Myrzakulov, Eur. Phys. J. C **71**, 1752 (2011);
P.Y. Tsyba, et al., Int. J. Theor. Phys. **50**, 1876 (2011).

[4] R. Ferraro, F. Fiorini, Phys. Rev. D **75**, 084031 (2007);
R. Ferraro, F. Fiorini, Phys. Rev. D **78**, 124019 (2008).

[5] P. Wu, H. Yu, Phys. Lett. B **692**, 176 (2010);
P. Wu, H. Yu, Eur. Phys. J. C **71**, 1552 (2011);
H. Wei, X.P. Ma, H.Y. Qi, Phys. Lett. B **703**, 74 (2011).

[6] P. Wu, H. Yu, Phys. Lett. B **693**, 415 (2010).

[7] K. Karami, A. Abdolmaleki, arXiv:1009.2459;
K. Karami, A. Abdolmaleki, arXiv:1009.3587;
K. Karami, A. Abdolmaleki, arXiv:1111.7269.

[8] E.V. Linder, Phys. Rev. D **81**, 127301 (2010).

[9] J.B. Dent, S. Dutta, E.N. Saridakis, Phys. Rev. D **83**, 023508 (2011);
J.B. Dent, S. Dutta, E.N. Saridakis, JCAP **01**, 009 (2011);
C.Q. Geng, C.C. Lee, E.N. Saridakis, Y.P. Wu, Phys. Lett. B **704**, 384 (2011);
C.Q. Geng, C.C. Lee, E.N. Saridakis, JCAP **01**, 002 (2012).

[10] M. Li, R.X. Miao, Y.G. Miao, JHEP **07**, 108 (2011).

[11] F.R. Urban, A.R. Zhitnitsky, Phys. Rev. D **80**, 063001 (2009);
F.R. Urban, A.R. Zhitnitsky, JCAP **09**, 018 (2009);
F.R. Urban, A.R. Zhitnitsky, Phys. Lett. B **688**, 9 (2010);
F.R. Urban, A.R. Zhitnitsky, Nucl. Phys. B **835**, 135 (2010);
N. Ohta, Phys. Lett. B **695**, 41 (2011);
R.G. Cai, Z.L. Tuo, H.B. Zhang, arXiv:1011.3212;
R.G. Cai, Z.L. Tuo, Y.B. Wu, Y.Y. Zhao, arXiv:1201.2494.

[12] E. Ebrahimi, A. Sheykhi, Phys. Lett. B **705**, 19 (2011);
E. Ebrahimi, A. Sheykhi, Int. J. Mod. Phys. D **20**, 2369 (2011);
A. Sheykhi, A. Bagheri, Europhys. Lett. **95**, 39001 (2011);
A. Sheykhi, M. Sadegh Movahed, Gen. Relativ. Gravit. **44**, 449 (2012);
A. Rozas-Fernandez, arXiv:1106.0056;
A. Khodam-Mohammadi, M. Malekjani, arXiv:1201.3200.

[13] S. Capozziello, V.F. Cardone, H. Farajollahi, A. Ravanpak, Phys. Rev. D **84**, 043527 (2011).

[14] R.R. Caldwell, E.V. Linder, Phys. Rev. Lett. **95**, 141301 (2005).

[15] E.E.O. Ishida, R.R.R. Reis, A.V. Toribio, I. Waga, Astropart. Phys. **28**, 547 (2008).



Figure 1: The evolution of ghost $f(T)$-gravity model, Eq. (23), versus T/T_0. The auxiliary parameters are $H_0 = 71.8$ Km S^{-1} Mpc^{-1} and $\Omega_{m_0} = 0.26$ [13]. For these values one finds $\beta = 130.5$.



Figure 2: The evolution of the torsion scaler, Eq. (27), versus t/t_0. Auxiliary parameters as in Fig. 1.



Figure 3: The evolution of the effective torsion EoS parameter of the ghost $f(T)$-gravity model, Eq. (28), versus t/t_0. Auxiliary parameters as in Fig. 1.



Figure 4: The evolution of the deceleration parameter of the ghost $f(T)$-gravity model, Eq. (29), versus t/t_0. Auxiliary parameters as in Fig. 1.